

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 6 : C12N 9/00		A2	(11) International Publication Number: WO 97/09420 (43) International Publication Date: 13 March 1997 (13.03.97)
(21) International Application Number: PCT/GB96/02180 (22) International Filing Date: 5 September 1996 (05.09.96) (30) Priority Data: 9518026.1 5 September 1995 (05.09.95) GB 9521496.1 20 October 1995 (20.10.95) GB 9521497.9 20 October 1995 (20.10.95) GB 9526230.9 21 December 1995 (21.12.95) GB 9612145.4 11 June 1996 (11.06.96) GB		(81) Designated States: AL, AM, AT, AU, AZ, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, US, UZ, VN, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>Without international search report and to be republished upon receipt of that report.</i>	
(71) Applicant (for all designated States except US): CELLTECH THERAPEUTICS LIMITED [GB/GB]; 216 Bath Road, Slough, Berkshire SL1 4EN (GB). (72) Inventors; and (75) Inventors/Applicants (for US only): DOCHERTY, Andrew, James, Penrose [GB/GB]; Piccards Cottage, Sandy Lane, Guildford, Surrey GU3 1HF (GB). SLOCOMBE, Patrick, Marcel [GB/GB]; 16 Roman Way, Warfield, Bracknell, Berkshire RG42 7UT (GB). (74) Agent: HALLYBONE, Huw, George; Carpmaels & Ransford, 43 Bloomsbury Square, London, WC1A 2RA (GB).			
(54) Title: DNA SEQUENCES CODING FOR A HUMAN METALLOPROTEINASE AND VARIANTS THEREOF			
(57) Abstract DNA sequences coding for a human metalloproteinase are described together with the corresponding antisense DNA and RNA. The DNA may be used to produce the metalloproteinase which may be used to generate antibodies thereto and to obtain other compounds capable of regulating the action of the metalloproteinase <i>in vivo</i> .			

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Larvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

DNA SEQUENCES CODING FOR A HUMAN
METALLOPROTEINASE AND VARIANTS THEREOF

5 This invention relates to a novel human metalloproteinase, to homologues and fragments thereof, to means for producing the metalloproteinases, and to means for regulating their production and activity *in vivo*.

A number of physiologically important processing events are mediated by 10 metalloproteinases, which under certain circumstances contribute to pathologies as diverse as inflammation and cancer, and it has been suggested that such enzymes would provide targets for therapeutic intervention. Thus, by varying the production of the enzyme, or inhibiting or enhancing its activity *in vivo* it should be possible to achieve a 15 therapeutic effect.

In one example, tumour necrosis factor-alpha (TNF- α) is a potent pro-inflammatory and immunomodulatory mammalian cytokine produced primarily by activated monocytes and macrophages. It is initially 20 expressed as a 233-amino-acid membrane-anchored precursor (pro-TNF- α) which is proteolytically processed to yield the mature, 157-amino-acid cytokine. Evidence has been obtained which indicates that at least one metalloproteinase-like enzyme mediates pro-TNF- α cleavage, but to date the enzyme(s) responsible for this *in vivo* are unknown [see for example 25 Mohler, K M *et al*, *Nature* 370, 218-220 (1994); Gearing, A J *et al* *ibid* 370, 555-557 (1994); McGeehan, G M *et al*, *ibid* 370, 558-561 (1994)]. A number of known matrix metalloproteinase inhibitors have been shown to block TNF- α secretion [see the above papers and International Patent Specification Publication No. WO 95/06031]. These compounds were 30 originally designed to selectively inhibit matrix metalloproteinases such as collagenase with primary functions unrelated to pro-TNF- α cleavage. Where new inhibitors have been described these have apparently been selected on the basis of their effect on TNF- α secretion seen in cell-based assays.

In another example, L-selectin shedding is thought to be a pro-inflammatory event that is mediated by an as yet unidentified metalloproteinase [Lasky, *Science*, 258, 964-969 (1992)]. Some inhibitors of L-selectin proteolysis have been identified, but these have been 5 obtained using cell based assays [Walchech *et al.*, *Nature*, 380, 720-723 (1996); Feehan *et al.*, *J. Biol. Chem.*, 271, 7019-7024 (1996)].

In general, in order to obtain compounds capable of selectively regulating 10 the action of a metalloproteinase implicated in human disease, for example as in the above TNF- α and L-selectin instances it would be clearly advantageous to have the enzyme unequivocally identified and obtainable in an isolated, purified and unambiguous characterised form.

Through the use of a cloning and screening approach, we have been able 15 to identify human DNA which is responsible for coding part of one such metalloproteinase. This DNA has the sequence described in SEQ I.D. No: 1 below and may be of use (1) in the generation of a gene coding for the metalloproteinase, (2) in the production of the metalloproteinase, (3) in the provision of means to regulate the activity of the metalloproteinase *in vivo*, 20 and (4) in the provision of means to detect and measure a metalloproteinase in a biological system, e.g. in serum, synovial fluid or a tissue extract.

Thus according to one aspect of the invention we provide DNA comprising 25 the nucleotide sequence of SEQ I.D. No: 1:

SEQ I.D. No: 1

GGTCGAGCACTCCAAGCCCACCACCAAGGGACTGGGCTCTCAGTT
30 ACACAACAGACCAAGAAGCGACCTCGCAGGATGAAAAGGAAAGATT
AAACTCCATGAAGTATGTGGAGCTTACCTCGTGGCTGATTATTAGA
GTTTCAGAAGAATCGACGAGACCAGGACGCCACAAACACAAGCTCA
TAGAGATGCCAACTATGTTGATAAGTTTACCGATCCTTGAACATCC
GGATTGCTCTCGTGGCTTGGAAAGTGTGGACCCACGGAACATGTG
35 TGAAGTTTCAGAGAATCCATATTCTACCCTCTGGTCTTCTCAGTTG
GAGGCGCAAGCTGCTGCCAGAAGTACCATGACAACGCCAATTAA

TCACGGGCATGTCCTCCACGGCACCAACCACATGGCCTGGCCCCCT
CATGGCCATGTGCTCTGTGTACCAAGTCTGGAGGAAGTCAACATGGACC
ACTCCGAGAATGCCATTGGCGTGGCTGCCACCATGGCCCACGAGAT
GGGCCACAACTTGGCATGACCCATGATTCTGCAGATTGCTGCTCGG
5 CCAGTGCAGGCTGATGGTGGGTGCATCATGGCAGCTGCCACTGGGCA
CCCCTTCCCAAAGTGTCAATGGATGCAACAGGAGGGAGCTGGACA
GGTATCTGCAGTCAGGTGGATGTGTCTCTCCAACATGCCAGAC
ACCAGGATGTTGTATGGAGGCCGGAGGTGTGGGAACGGGTATCTGG
AAGATGGGAAGAGTGTGACTGTGGAGAAGAAGAGGAATGTAACAA
10 CCCCTGCTGCAATGCCTCTAATTGTACCCCTGAGGCCGGGGCGGAG
TGTGCTCACGGCTCCTGCTGCCACCAAGTGTAAAGCTGTTGGCTCGTGG
GACCCCTGTGCCCGAGCAGGCCAGGCAGTGTGACCTCCCGGAGTTC
TGTACGGGCAAGTCTCCCCACTGCCCTACCAACTTCTACCAGATGGA
TGGTACCCCTGTGAGGGCGGCCAGGCCTACTGCTACAAACGGCATG
15 TGCCTCACCTACCAGGAGCAGTGCAGCAGCTGTGGGGACCCGGAG
CCCGACCTGCCCTGACCTCTGCTCGAGAAGGTGAATGTGGCAGG
AGACACCTTGGAAACTGTGGAAAGGACATGAATGGTAACACAGGA
AGTGCAACATGAGAGATGCGAAGTGTGGGAAGATCCAGTGTCAAG
CTCTGAGGCCGGCCCTGGAGTCCAACGCCGTGCCATTGACACC
20 ACTATCATCATGAATGGGAGGCAGATCCAGTGCAGCAGCTGTGGGGACCCACG
TCTACCGAGGTCTGAGGAGGGAGGGTGACATGCTGGACCCAGGGCT
GGTGAATGACTGGAACCAAGTGTGGCTACAACCATAATTGCTTGAGG
GGCAGTGCAGGAACACCTCCCTTTGAAACTGAAGGCTGTGGGAAG
AAGTGCAATGCCATGGGCTGTAAACAACAACCAGAACTGCCACTG
25 CCTGCCGGCTGGCCCCGCCCTTGCAACACACCCGGGCCACGG
GGGCAGTATCGACAGTGGCCTATGCCCTGAGAGTGTGGTCCT
GTGGTAGCTGGAGTGTGGCTGGCCATCTGGTGCTGGCGGTCTCA
TGCTGATGTACTACTGCTGCAGACAGAACAAACTAGGCCAACTC
AAGCCCTCAGCTCTCCCTCCAAGCTGAGGCAACAGTTCAGTTGTCC
30 CTTCAGGGTTCTCAGAACAGCGGGACTGGTCATGCCAACCCAACTT
TCAAGCTGCAGACGCCAGGGCAAGCGAAAGGTGTTCTTGACTT
GTGCGTACAGGTGATCAACACTCCGGAAATCCTGCGGAAGCCCTCC
CAGCCTCCCTCCCCGGCCCCCTCCAGATTATCTGCGTGGTGGTCCC
CACCTGCACCACTGCCAGCTCACCTGAGCAGGGCTGCTAGGAAC
35 CCCAGGGCCGGGTCTCAAATAGAGAGGACGGAGTCGTCCAGGAG
GCCTCCTCCAAGCCGCCAATTCCCCCGCACCAAATTGCATCGTTT

CCCAGGACTTCTCCAGGCCTGGCCGCCAGAAGGCACCTCCGGC
AAACCCAGTGCAGGCCGCAGGAGCCTCCCCAGGCCAGGAGGTGC
ATCCCCACTGCAGGCCCTGGTGCCTGGCCCTCAGCAGTCCGGCCT
CTGGCAGCACTTGCCCCAAAGAGGGTATGGAAGACTTGCAATTGAA
5 AACTGGGGACCAGTCCAAAGTCAGTAATTGTGTTAACACGTGTATA
ACAGCTCTGCTGGACACCCAAGAAAGCCATGGGAACGCCAAGTGG
AAGGTCCCCCTCCCCAGGGAGCCTGCGAAGGAGAGGTTCTGTAGA
ATCCAAGCCCACATTCCAAAGTCACCCCCAACGCGTCCTCTCACAC
CGTCCACTGTGCCTTGTATGTGTCAGGGATCCAGGGCAATGTGAAT
10 TTTCTTTTATTGGGAGATTGTTACGGAAAACAGATCTCTCTCTC
TTGTCCACCTATTAAATTGTTACAATATTGTACATCTATGCAAAATAC
TTGAATGGGCCATGGTGCCTTTCTCTGTTAGTATTAAATTAAAAAAT
GAATTGTTGTCATTGCAAAAAAAAAAAAAAA

15 and homologues and fragments thereof.

It will be appreciated that the nucleotide sequence of SEQ I.D. No: 1 also includes control sequences, such as a polyadenylation sequence, providing for expression of the sequence in a host cell.

20 One particular DNA fragment according to the invention is the isolated human metalloproteinase-encoding nucleotide sequence of SEQ I.D. No: 2:

25 **SEQ I.D. No: 2**

TTCGAGCACTCCAAGCCCACCACCAAGGGACTGGCTTCAAGTTAC
ACAACAGACCAAGAACGACCTCGCAGGATGAAAAGGGAAAGATTAA
ACTCCATGAAGTATGTGGAGCTTACCTCGTGGCTGATTATTAGAGT
30 TTCAGAAGAACATCGACGAGACCAGGACGCCACAAACACAAGCTCATA
GAGATGCCAACTATGTTGATAAGTTTACCGATCCTGAACATCCGG
ATTGCTCTCGTGGCTTGGAAAGTGTGGACCCACGGGAACATGTGTG
AAGTTTCAGAGAACATCCATATTCTACCCCTCTGGCCTTCTCAGTTGGA
GGCGCAAGCTGCTGCCAGAAGTACCATGACAACGCCAATTATC
35 ACGGGCATGTCCTTCCACGGCACCCATCGGCCTGGCCCCCCTCA
TGGCCATGTGCTCTGTGTACCAAGTCTGGAGGAGTCAACATGGACAC

TCCGAGAATGCCATTGGCGTGGCTGCCACCATGGCCCACGAGATGG
GCCACAACTTGGCATGACCCATGATTCTGCAGATTGCTGCTCGGCC
AGTGCAGCTGATGGTGGGTGCATCATGGCAGCTGCCACTGGCACC
CCTTCCCAAAGTGTCAATGGATGCAACAGGAGGGAGCTGGACAG
5 GTATCTGCAGTCAGGTGGATGTGTCTCTCCAACATGCCAGACA
CCAGGATGTTGTATGGAGGCCGGAGGTGTGGAACGGGTATCTGGA
AGATGGGAAGAGTGTGACTGTGGAGAAGAAGAGGAATGTAACAAC
CCCTGCTGCAATGCCTCTAATTGTACCCCTGAGGCCGGGGCGGAGT
GTGCTCACGGCTCCTGCTGCCACCAAGTGTAAAGCTTGGCTCCTGG
10 GACCCTGTGCCCGCAGCAGGCCAGGCAGTGTGACCTCCGGAGTTC
TGTACGGGCAAGTCTCCCCACTGCCCTACCAACTTCTACCAGATGGA
TGGTACCCCCCTGTGAGGGCGGCCAGGCCTACTGCTACAAACGGCATG
TGCCTCACCTACCAGGAGCAGTGCAGCAGCTGTGGGGACCCGGAG
CCCGACCTGCCCTGACCTCTGCTCGAGAAGGTGAATGTGGCAGG
15 GACACCTTGGAAACTGTGGAAAGGACATGAATGGTGAACACAGGAA
GTGCAACATGAGAGATGCGAAGTGTGGAAAGATCCAGTGTCAAGGC
TCTGAGGCCGGCCCTGGAGTCCAACCGGGTGCCATTGACACCA
CTATCATCATGAATGGGAGGCAGATCCAGTGCCGGGGACCCACGT
CTACCGAGGTCTGAGGAGGGAGGGTACATGCTGGACCCAGGGCTG
20 GTGATGACTGGAACCAAGTGTGGCTACAACCATAATTGCTTGAGGG
CAGTGCAGGAACACCTCCTCTTGAAACTGAAGGCTGTGGGAAGAA
GTGCAATGGCCATGGGTCTGTAACAACAACCAGAACTGCCACTGCC
TGCCGGCTGGGCCCTGCAACACACCGGGCACGGGG
GCAGTATCGACAGTGGGCCTATGCCCTGAGAGTGTGGGTCTGT
25 GGTAGCTGGAGTGTGGGCCATCTTGGTGTGGCTGGCGGTCTCATG
CTGATGTACTACTGCTGCAGACAGAACAAACTAGGCCAACTCAA
GCCCTCAGCTCTCCCTCCAAGCTGAGGCAACAGTTCAAGTGTCCCT
TCAGGGTTCTCAGAACAGCGGGACTGGTCATGCCAACCAACTTTC
AAGCTGCAGACGCCAGGGCAAGCGAAAGGTGTCCCTGACTTGT
30 GCGTACAGGTGATCAACACTCCGGAAATCCTGCAGGAAAGCCCTCCCA
GCCTCCTCCCCGGCCCCCTCCAGATTCTGCGTGGTGGTCCCT
CCTGCACCACTGCCAGCTCACCTGAGCAGGGCTGCTAGGAACCTCC
CAGGGCCCGGGTCTCAAATAGAGAGGACGGAGTCGTCCAGGAGGC
CTCCCTCCAAGGCCAGGGCAATTCCCCCGCACCAAATTGCATCGTTCC
35 CAGGACTTCTCCAGGCCTGGCCGGCCCCAGAAGGCAGTCCAGGAGGTGCAT
ACCCAGTGCCAGGCCAGGGCAGGAGCCTCCCCAGGCCAGGAGGTGCAT

CCCCACTGCGGCCCTGGTGCTGGCCCTCAGCAGTCCCAGCCTCT
GGCAGCACCTGCCAAAGAGGGTATGGAAGACTTGCAATTGAAAAA
CTGGGGACCAGTTCCAAAGTCAG

5 and homologues and fragments thereof.

In the sequences herein standard one letter codes are used to represent nucleotides or amino acids as appropriate.

10 DNA according to the invention may be obtained using conventional molecular biology procedures, for example by probing a human genomic or cDNA library with one or more labelled oligonucleotide probes containing for example fifteen or more contiguous nucleotides designed using the nucleotide sequences described herein [see for example 15 "Current Protocols in Molecular Biology", Ausubel, F M *et al* (eds), Greene Publishing Associates and Wiley-Interscience, New York (1987)].

Where the term homologue is used herein in relation to a particular nucleotide or amino acid sequence this is to be understood to represent a 20 corresponding sequence in which one or more nucleotides or amino acids have been added, deleted, substituted or otherwise chemically modified, provided always that the homologue retains substantially the same catalytic properties as the particular sequence described. One particular type of homologue for example may be that in which one or more 25 nucleotides have been substituted due to the degeneracy of the genetic code. Homologues, particularly longer versions of the sequences described herein, may be obtained by standard molecular biology and/or chemistry techniques, e.g. by cDNA or gene cloning, or by use of oligonucleotide directed mutagenesis or oligonucleotide directed synthesis 30 techniques or enzymatic cleavage or enzymatic filling in of gapped oligonucleotides (see for example Ausubel, F M *ibid*).

The DNA of SEQ I.D. No: 1 and SEQ I.D. No: 2 each codes for part of a 35 human metalloproteinase. Thus, the DNA according to the invention or a fragment thereof may be used as a probe to screen an appropriate genomic or cDNA library in a process utilising standard hybridisation

and/or PCR cloning techniques to obtain the gene or cDNA coding for the entire metalloproteinase, or a homologue or fragment thereof, or a related metalloproteinase from another species.

- 5 Thus according to a further aspect of the invention we provide an isolated gene or cDNA coding for a human metalloproteinase, said gene or cDNA containing the nucleotide sequence of SEQ I.D. No: 1 or SEQ I.D. No: 2 or a homologue thereof.
- 10 The gene according to the invention may in turn be used to produce a metalloproteinase. In another aspect of the invention we therefore provide an isolated human metalloproteinase which contains the amino acid sequence of SEQ I.D. No: 3:

15

SEQ I.D. No: 3

FEHSKPTTRDWALQFTQQTKRPRRMKREDLNSMKYVELYLVADYLEF
QKNRRDQDATKHKLIEIANVYDKFYRSLNIRIALVGEVWTHGNMCEVSE
NPYSTLWSFLSWRRKLLAQKYHDNAQLITGMSFHGTTIGLAPLMAMCSV
20 YQSGGVNMDHSENAIGVAATMAHEMGHNGMTHDSADCCSASAADG
GCIMAAATGHPFPKVFNCGNRRELDRLQSGGGMCLSNMPDTRMLYG
GRRCGNGYLEDGECCDCGEEEECNPCCNASNCTLRPGAECAHGS
HQCKLLAPGTLCREQARQCDLPEFCTGKSPHCPTNFYQMDGTPCEGG
QAYCYNGMCLTYQEQCQQLWGPGPARPAPDLCFEKVNAGDTFGNC
25 KDMNGEHRKCNMRDAKCGKIQCQSSEARPLESNAVPIDTTIIMNGRQIQ
CRGTHVYRGPEEEGDMLDPGLVMTGKCGYNHICFEGQCRNTSFFETE
GCGKKCNGHGVCNNNNQNCHCLPGWAPPFCNTPGHGGSIDSGPMPPE
SVPVVAGVLVAILVLAVLMLMYYCCRQNNKLGQLKPSALPSKLRQQFS
CPFRVSQNSGTGHANPTFKLQTPQGKRKVFLDCVQVINTPEILRKPSQ
30 PPPRPPPDYLRGGSPPAPLPAHLSRAARNSPGPGSQIERTESSRRPPPS
RPIPPAPNCIVSQDFSRPRPPQKALPANPVPGRRLSPRGGASPLRPPG
AGPQQSRPLAALAPKRVWKTCLKTGDQFQSQ

and homologues and fragments thereof.

35

The production of a protein according to the invention may be achieved using standard recombinant DNA techniques involving the expression of the metalloproteinase by a host cell. The isolated nucleic acids described herein may be for example introduced into any suitable expression vector

5 by operatively linking the DNA to any necessary expression control elements therein and transforming any suitable prokaryotic or eucaryotic host cell with the vector using well known procedures. The invention is thus to be understood to extend to recombinant plasmids containing a gene of the invention or a nucleotide sequence of SEQ I.D. No: 1 or SEQ

10 I.D. No: 2, to cells containing said recombinant plasmids and to a process for producing the protein according to the invention which comprises culturing said cells such that the desired protein is expressed and recovering the protein from the culture.

15 Thus in one example the nucleotide sequence of SEQ I.D. No: 1, without its 3' poly A tail, or SEQ I.D. No: 2, or a homologue such as a longer version including a sequence encoding a signal peptide for secretion, and a propeptide to ensure accurate enzyme folding, is inserted downstream of the hCMV promoter in the pEE12 plasmid vector, and either transiently

20 or stably expressed in CHO-L761h or NSO mouse melanoma cells [Murphy et al., J. Biol. Chem., 267, 9612-9618 (1992)]. Expression of the enzyme according to the invention can be detected in serum free culture medium by its catalytic properties, or by Western blotting [Murphy et al., Biochem. J., 283, 637-641 (1992); Murphy et al., J. Biol. Chem., 267, 9612-9618 (1992)]. Such assays can also be used during the subsequent

25 isolation of the expressed enzyme from transfected cell conditioned medium. If the enzyme requires further activation, this may be achieved proteolytically through use of modest amounts of trypsin, furin, or other methods, in order to remove an approximately 180 amino acid N-terminal propeptide, as described for other metalloproteinases [Murphy et al., J. Biol. Chem., 267, 9612-9618 (1992); Crabbe et al., Biochemistry, 33, 14419-14425 (1994); Pei and Weiss, Nature, 375, 244-247 (1995); Will et al., J. Biol. Chem., 271, 17119-17123 (1996)]

35 It may be desirable to produce the catalytic domain of the protein according to the invention in isolation from the rest of the molecule. This

may be achieved using the above standard recombinant DNA techniques except that in this instance the DNA sequence used is that encoding the amino acid sequence of SEQ I.D. No: 4:

5

SEQ I.D. No: 4

MKREDLNSMKYVELYLVADYLEFQKNRRDQDATKHKLIEIANYVDKFYR
SLNIRIALVGLEVWTHGNMCEVSENPYSTLWSFLSWRRKLLAQKYHDNA
QLITGMSFHGTTIGLAPLMAMCSVYQSGGVNMDHSENAIGVAATMAHE
10 MGHNFGMTHDSADCCSASAADGGCIMAAATGHPFPKVNGCNRRELD
RYLQSGGGMCLSNMPDTRMLYG

or a homologue thereof, and the invention extends to such isolated catalytic domains.

15

N or C-terminally extended versions of the sequence shown in SEQ I.D. No: 4 may be obtained by expression in prokaryotic or eucaryotic cells as described above optionally attached to a peptide tag via which the protein may be affinity purified and identified. Examples of tags include the well known "His" or "Strep-tags". Further sequences that may be attached arise from expression in prokaryotic cells and include the *pelB* or *ompA* leaders which when placed at the N-terminus help direct secretion to the *E.coli* periplasmic space [Schmidt and Skerra, *J. Chromatography*, 676, 337-345 (1994); Knauper et al., *J. Biol. Chem.*, 271, 17124-17131 (1996)].

The gene or nucleotide sequences according to the invention may also be of use in diagnosis, for example to determine enzyme deficiency in a human subject, by for example direct DNA sequence comparison or 30 DNA/RNA hybridisation assays; or in therapy, for example where it is desired to modify the production of the metalloproteinase *in vivo*, and the invention extends to such uses.

Knowledge of the gene according to the invention also provides the ability 35 to regulate its activity *in vivo* by for example the use of antisense DNA or RNA. Thus, according to a further aspect of the invention we provide an

antisense DNA or an antisense RNA of a gene coding for a human metalloproteinase, said gene containing the nucleotide sequence of formulae SEQ I.D. No: 1 or SEQ I.D. No: 2.

- 5 The antisense DNA or RNA will correspond to the metalloproteinase gene or a fragment thereof, for example a fragment based on the nucleotide sequence of SEQ I.D. No: 1 or SEQ I.D. No: 2. The antisense DNA or RNA can be produced using conventional means, by standard molecular biology and/or by chemical synthesis as described above. If desired, the
- 10 antisense DNA and antisense RNA may be chemically modified so as to prevent degradation *in vivo* or to facilitate passage through a cell membrane, and/or a substance capable of inactivating mRNA, for example ribosyme, may be linked thereto, and the invention extends to such constructs.
- 15 The antisense DNA or antisense RNA may be of use in the treatment of diseases or disorders in humans in which the over- or unregulated production of the metalloproteinase has been implicated. Such diseases or disorders may include those described under the general headings of
- 20 infectious diseases, e.g. HIV infection; inflammatory disease/autoimmunity e.g. rheumatoid arthritis, inflammatory bowel disease; osteoarthritis; cancer; allergic/atopic diseases e.g. asthma, eczema; cardiovascular disease e.g. myocardial infarction, congestive heart failure; systemic inflammatory response syndrome e.g. sepsis syndrome; reperfusion injury;
- 25 malignancy; cachexia; congenital e.g. cystic fibrosis, sickle cell anaemia; dermatologic, e.g. psoriasis, alopecia; neurologic, e.g. multiple sclerosis, migraine headache; renal e.g. uraemia, nephrotic syndrome; obstetric/gynecologic e.g. premature labour, miscarriage, genitourinary prolapse, urinary incontinence, contraception, infertility; transplants e.g.
- 30 organ transplant rejection, graft-versus-host disease; metabolic/idiopathic disease e.g. diabetes; disorders of the bone such as osteoporosis; and toxicity e.g. due to chemotherapy, cytokine therapy, and anti-CD3 therapy.

The metalloproteinase according to the invention and homologues or fragments thereof may be used to generate substances which selectively bind to the proteins and in so doing regulate the activity of the enzymes.

Such substances include, for example, antibodies, and the invention extends in particular to an antibody which is capable of recognising one or more epitopes on a metalloproteinase containing the amino acid sequence of SEQ I.D. No: 3, or a homologue or fragment thereof. In particular the 5 antibody may be a neutralising antibody

As used herein the term antibody is to be understood to mean a whole antibody or a fragment thereof, for example a F(ab)₂, Fab, Fv, V_H or V_K fragment, a single-chain antibody, a multimeric monospecific antibody or 10 fragment thereof, or a bi- or multispecific antibody or fragment thereof.

The antibody according to the invention may be a polyclonal or, especially, a monoclonal antibody. The antibody may belong to any immunoglobulin class, and may be for example an IgG, for example IgG₁, IgG₂, IgG₃ 15 IgG₄, IgE, IgM or IgA antibody. It may be of animal, for example mammalian origin, and may be for example a murine, rat or human antibody. Alternatively, the antibody may be a chimeric antibody. The term chimeric antibody is used herein to mean any antibody containing portions derived from different animal species. Particular examples 20 include those antibodies having a variable region derived from a murine or other antibody constant region, and those antibodies in which one or more CDR sequences and optionally one or more variable region framework amino acids are derived from a murine or other antibody and the remaining portions of the variable and the constant regions are derived 25 from a human immunoglobulin.

Antibodies according to the invention may be prepared by conventional immunisation and recombinant DNA techniques. Thus, for example polyclonal antibodies may be obtained from the sera of animals 30 immunised with a metalloproteinase according to the invention or a homologue or fragment thereof. Any suitable host, for example BALB/c mice where it is desired to obtain a mouse polyclonal antibody, may be injected with the immunogen, the serum collected and the antibody recovered therefrom. Monoclonal antibodies may be obtained from 35 hybridomas derived from the spleen cells of an animal immunised as just discussed and fused to an appropriate "immortal" B-tumour cell. In each

instance, the antibody may be recovered from either the serum or the hybridoma by making use of standard purification and/or concentration techniques, for example by chromatography, using for example Protein A or by other affinity chromatography employing a metalloproteinase of the 5 invention or a homologue or fragment thereof.

Once a cell line, for example a hybridoma, expressing an antibody according to the invention has been obtained it is possible to clone therefrom the cDNA and to identify the variable region genes encoding the 10 desired antibody, including the sequences encoding the CDRs. From here, other chimeric antibodies according to the invention may be obtained by preparing one or more replicable expression vectors containing at least the DNA sequence encoding the variable domain of the antibody heavy or light chain and optionally other DNA sequences encoding remaining 15 portions of the heavy and/or light chains as desired, and transforming an appropriate cell line, e.g. a non-producing myeloma cell line, such as a mouse NSO line, in which production of the antibody will occur. In order to obtain efficient transcription and translation, the DNA sequence in each vector should include appropriate regulatory sequences, particularly a 20 promoter and leader sequence operably linked to the variable domain sequence. Particular methods for producing antibodies in this way are generally well known and routinely used. For example, basic molecular biology procedures are described by Maniatis *et al* [Molecular Cloning, Cold Spring Harbor Laboratory, New York, 1989]; DNA sequencing can be 25 performed as described in Sanger *et al* [PNAS 74, 5463, (1977)] and the Amersham International plc sequencing handbook; and site directed mutagenesis can be carried out according to the method of Kramer *et al* [Nucl. Acids Res. 12, 9441, (1984)] and the Anglian Biotechnology Ltd handbook. Additionally, there are numerous publications, including patent 30 specifications, detailing techniques suitable for the preparation of antibodies by manipulation of DNA, creation of expression vectors and transformation of appropriate cells, for example as reviewed by Mountain A and Adair, J R in Biotechnology and Genetic Engineering Reviews [ed. Tombs, M P, 10, Chapter 1, 1992, Intercept, Andover, UK] and in 35 International Patent Specification No. WO 91/09967.

Antibodies and other selective binding agents according to the invention may be of use in therapy, either alone or as a delivery agent, for the delivery of a drug, prodrug, radiometal or radioisotope for example in the treatment of diseases such as those described above in humans and/or

5 other animals, or may find a use as purification agents in the preparation of the human metalloproteinase or homologues or fragments thereof.

In a further use according to the invention, selective binding agents of the invention, such as antibodies, may form the basis of a diagnostic assay to

10 detect the presence or absence in a biological sample (e.g. serum, synovial fluid or a tissue extract) of a metalloproteinase as described herein. Thus for example the binding agent may be brought into contact with a sample of serum, synovial fluid or tissue under conditions in which a complex is formed between the binding agent and target

15 metalloproteinase. Qualitative and/or quantitative detection of the complex can then be used to determine the presence or absence of the metalloproteinase and in particular whether the enzyme is present in an abnormal quantity associated for example with a particular disease state.

20 The metalloproteinases according to the invention may in particular be used to screen for compounds which regulate the activity of the enzymes and the invention extends to such a screen and to the use of compounds obtainable therefrom to regulate metalloproteinases *in vivo*.

25 Thus according to a further aspect of the invention we provide a process for obtaining a compound capable of regulating the action of a human metalloproteinase *in vivo* which comprises subjecting one or more test compounds to a screen comprising (A) a metalloproteinase containing the amino acid sequence of SEQ I.D. No: 3 or a homologue or fragment thereof, or (B) a host cell transformed to be capable of expressing a metalloproteinase gene or cDNA or a homologue or fragment thereof containing a nucleotide sequence of SEQ I.D. No: 1 or SEQ I.D. No: 2 or a homologue or fragment thereof.

30

35 The screen according to the invention may be operated using conventional procedures, for example by bringing the test compound or compounds to

be screened and an appropriate substrate into contact with the metalloproteinase or a cell capable of producing it and determining affinity for the protein in accordance with standard practice.

- 5 Any compound obtainable in this way may have a potential use in the treatment in humans and/or other animals of one or more of the above mentioned diseases. The invention thus extends to a compound selected through its ability to regulate the activity of a metalloproteinase *in vivo* as primarily determined in a screening assay utilising a metalloproteinase
- 10 containing an amino acid sequence of SEQ I.D. No: 3 or a homologue or fragment thereof or a gene coding therefor, for use in the treatment of a disease in which the over- or under-activity or unregulated activity of the metalloproteinase is implicated.

CLAIMS

5

1. Isolated human DNA comprising the nucleotide sequence of SEQ I.D. No: 1 herein and homologues and fragments thereof.
- 10 2. Isolated DNA according to Claim 1 having the human metalloproteinase-encoding nucleotide sequence of SEQ I.D. No: 2 herein and homologues and fragments thereof.
- 15 3. An isolated gene or cDNA coding for a human metalloproteinase, said gene or cDNA containing the nucleotide sequence of SEQ I.D. No: 1 or SEQ I.D. No: 2 herein or a homologue thereof.
- 20 4. An antisense DNA or an antisense RNA of a gene coding for a human metalloproteinase, said gene containing the nucleotide sequence of SEQ I.D. No: 1 or SEQ I.D. No: 2 herein.
- 25 5. An isolated human metalloproteinase which contains the amino acid sequence of SEQ I.D. No: 3 and homologues and fragments thereof.
6. An antibody which is capable of recognising one or more epitopes of a metalloproteinase according to Claim 5 or a homologue or fragment thereof.
- 30 7. A process for obtaining a compound capable of regulating the action of a human metalloproteinase *in vivo* which comprises subjecting one or more compounds to a screen comprising a metalloproteinase according to Claim 5 or a homologue or fragment thereof.

8. A process for obtaining a compound capable of regulating a human metalloproteinase *in vivo* which comprises subjecting one or more test compounds to a screen comprising a host cell transformed to be capable of expressing a metalloproteinase gene or cDNA containing a nucleotide sequence according to Claims 1 or 2 or a homologue thereof.
- 10 9. A compound selected through its ability to regulate the activity of a metalloproteinase *in vivo* as primarily determined in a screening assay utilising a metalloproteinase having an amino acid sequence of SEQ I.D. No: 3 or a homologue or fragment thereof or a gene coding therefor, for use in the treatment of a disease in which the over- or under-activity or unregulated activity of the metalloproteinase is implicated.

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12N 9/00		A3	(11) International Publication Number: WO 97/09420 (43) International Publication Date: 13 March 1997 (13.03.97)
(21) International Application Number: PCT/GB96/02180 (22) International Filing Date: 5 September 1996 (05.09.96)		(81) Designated States: AL, AM, AT, AU, AZ, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, US, UZ, VN, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).	
(30) Priority Data: 9518026.1 5 September 1995 (05.09.95) GB 9521496.1 20 October 1995 (20.10.95) GB 9521497.9 20 October 1995 (20.10.95) GB 9526230.9 21 December 1995 (21.12.95) GB 9612145.4 11 June 1996 (11.06.96) GB		Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>	
(71) Applicant (for all designated States except US): CELLTECH THERAPEUTICS LIMITED [GB/GB]; 216 Bath Road, Slough, Berkshire SL1 4EN (GB). (72) Inventors; and (75) Inventors/Applicants (for US only): DOCHERTY, Andrew, James, Penrose [GB/GB]; Piccards Cottage, Sandy Lane, Guildford, Surrey GU3 1HF (GB). SLOCOMBE, Patrick, Marcel [GB/GB]; 16 Roman Way, Warfield, Bracknell, Berkshire RG42 7UT (GB). (74) Agent: HALLYBONE, Huw, George; Carpmaels & Ransford, 43 Bloomsbury Square, London, WC1A 2RA (GB).		(88) Date of publication of the international search report: 24 April 1997 (24.04.97)	

(54) Title: DNA SEQUENCES CODING FOR A HUMAN METALLOPROTEINASE AND VARIANTS THEREOF

(57) Abstract

DNA sequences coding for a human metalloproteinase are described together with the corresponding antisense DNA and RNA. The DNA may be used to produce the metalloproteinase which may be used to generate antibodies thereto and to obtain other compounds capable of regulating the action of the metalloproteinase *in vivo*.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

INTERNATIONAL SEARCH REPORT

Intern. Appl. No.

PCT/GB 96/02180

A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 C12N9/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 6 C12N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	INTERNATIONAL IMMUNOLOGY, vol. 2, no. 6, 1990, OXFORD,GB, pages 585-591, XP000196854 S. YOSHIDA ET AL.: "Molecular cloning of cDNA encoding MS2 antigen, a novel cell surface antigen strongly expressed in murine monocytic lineage" see the whole document ---	1-3
A	WO 90 10062 A (BOARD OF REGENTS, THE UNIVERSITY OF TEXAS) 7 September 1990 see example 5 ---	1,6,8 -/-

Further documents are listed in the continuation of box C.

Patent family members are listed in annex.

* Special categories of cited documents :

- *'A' document defining the general state of the art which is not considered to be of particular relevance
- *'E' earlier document but published on or after the international filing date
- *'L' document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *'O' document referring to an oral disclosure, use, exhibition or other means
- *'P' document published prior to the international filing date but later than the priority date claimed

*'T' later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

*'X' document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

*'Y' document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

*'&' document member of the same patent family

2

Date of the actual completion of the international search

19 February 1997

Date of mailing of the international search report

10.03.97

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl.
Fax (+ 31-70) 340-3016

Authorized officer

Mateo Rosell, A.M.

INTERNATIONAL SEARCH REPORT

Intern. Application No.
PCT/GB 96/02180

C (Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 92 09701 A (INSTITUT NATIONALE DE LA SANTE ET DE LA RECHERCHE MEDICALE (INSERM)) 11 June 1992 see page 44, line 20-25 see example 6 see page 20, line 33 - page 23, line 6 ---	1,4,6,8, 9
A	WO 87 07907 A (CELL-TECH LIMITED) 30 December 1987 see page 6, line 32 - page 7, line 31 ---	6,9
A	TRENDS IN GENETICS, vol. 6, no. 4, 1990, OXFORD, GB, pages 121-125, XP000644304 L.M. MATRISIAN: "Metalloproteinases and their inhibitors in matrix remodeling" see the whole document ---	1-3
A	JOURNAL OF PHARMACEUTICAL SCIENCES, vol. 84, no. 4, 1 April 1995, COLUMBUS, US, pages 404-409, XP000617963 S.K. CHANDER ET AL.: "An in vivo model for screening peptidomimetic inhibitors of gelatinase A" see the whole document ---	7
P,X	NATURE, vol. 377, - 19 October 1995 LONDON, GB, pages 652-656, XP000196834 T. YAGAMI-HIROMASA ET AL.: "A metalloprotease-disintegrin participating in myoblast fusion" see the whole document, specially figure 1a ---	1-3,5,6
P,X	JOURNAL OF CELLULAR BIOLOGY, vol. 132, no. 4, 1996, NEW YORK, US, pages 717-726, XP000644308 G. WESKAMP ET AL.: "MDC9, a widely expressed cellular disintegrin containing cytoplasmic SH3 ligand domains" see the whole document ---	1-3
P,X	JOURNAL OF BIOLOGICAL CHEMISTRY, vol. 271, no. 9, 1996, MARYLAND, US, pages 4593-4596, XP000644305 J.R. KRAETZSCHMAR ET AL.: "Metarginin, a membrane-anchored metalloprotease-disintegrin protein with an RGD integrin binding sequence" see the whole document -----	1-3

INTERNATIONAL SEARCH REPORT

Information on patent family members

Intern. Application No

PCT/GB 96/02180

Patent document cited in search report	Publication date	Patent family member(s)		Publication date
WO-A-9010062	07-09-90	AU-B-	639906	12-08-93
		AU-A-	5104790	26-09-90
		CA-A-	2048695	28-09-90
		DE-D-	69005297	27-01-94
		DE-T-	69005297	21-04-94
		EP-A-	0460016	11-12-91
		ES-T-	2060146	16-11-94
		JP-T-	4505699	08-10-92
WO-A-9209701	11-06-92	GB-A-	2250993	24-06-92
		AU-B-	646914	10-03-94
		AU-A-	9089691	25-06-92
		CA-A-	2073860	22-05-92
		EP-A-	0513316	19-11-92
		HU-A-	65373	02-05-94
		NZ-A-	240688	27-09-94
		US-A-	5484726	16-01-96
		US-A-	5236844	17-08-93
WO-A-8707907	30-12-87	EP-A-	0269724	08-06-88
		JP-T-	63503514	22-12-88

